

Subobjects of the Successive Power Objects in the Topos $G - Set$

Apoloniusz Tyszk

Abstract. Let G be a group and let M be an object of the topos $G - Set$. We prove that an object X of the category $G - Set$ is isomorphic to some subobject of one of the objects $P(M), P(P(M)), P(P(P(M))), \dots$ if and only if $\text{card } X < \sup\{\text{card } P(M), \text{card } P(P(M)), \text{card } P(P(P(M))), \dots\}$ and $\{g \in G : \forall m \in M \ gm = m\} \subseteq \{g \in G : \forall x \in X \ gx = x\}$.

In this paper we present a simplified proof of a theorem which was proved in [6]; earlier weaker results of this type can be found in [4] and [5]. Let G be a group. The category $G - Set$ has as objects, sets equipped with a G -action and as morphisms functions preserving this action. This is a topos (see [2] for the introduction to the topos theory), i.e. $G - Set$ is an elementary topos in the terminology of [3]. Let $P : G - Set \rightarrow G - Set$ denote the power functor.

Theorem. *Let M be an object of the topos $G - Set$. Then an object X of the topos $G - Set$ is isomorphic to some subobject of one of the objects $M_n := \underbrace{P(\dots(P(M)))}_{n \text{ times}}$*

($n \geq 1$) if and only if

$$\text{card } X < \chi(M) := \sup\{\text{card } M, \text{card } P(M), \text{card } P(P(M)), \dots\}, \quad (1)$$

and

$$G_M := \{g \in G : \forall m \in M \ gm = m\} \subseteq G_X := \{g \in G : \forall x \in X \ gx = x\}. \quad (2)$$

Proof. The necessity is obvious, we show the sufficiency. Each transitive G -set X which satisfies (2) is isomorphic to a G -set of the form $(\{gH : g \in G\}, G, L)$ where H is a subgroup of G and L denotes left translation (see [1] p.106), obviously $G_M \subseteq G_X \subseteq H$. Take any relation which well orders M , let $M = \{m_\gamma\}_{\gamma < \alpha}$. Then the transitive G -set determined by $z := \{\{m_\gamma : \gamma < \beta\} : \beta \leq \alpha\} \in M_2$ is isomorphic to $(\{gG_M : g \in G\}, G, L)$, hence (Z. Moszner in [6]) $(\{gHz : g \in G\}, G, L)$ i.e. $(\{\{ghz : h \in H\} : g \in G\}, G, L)$ is a subobject of M_3 which is isomorphic to $(\{gH : g \in G\}, G, L)$.

An arbitrary G -set X which satisfies (1) and (2) is a disjoint union of less than $\chi(M)$ transitive G -sets Y_δ such that $G_M \subseteq G_{Y_\delta} := \{g \in G : \forall y \in Y_\delta \ gy = y\}$. The proof will be completed by showing that M_{n+4} includes $\text{card } M_n$ pairwise disjoint objects, each of them isomorphic to M_3 . Let $\text{Bij}(M)$ denote the group of bijections

of M , by $\mathcal{P} : Set \rightarrow Set$ we mean the power set functor. The relation:

$$xR_{n+1}y : \Longleftrightarrow \exists f \in Bij(M) \ y = \underbrace{\mathcal{P}(\dots(\mathcal{P}(f)))}_{n+1 \text{ times}}(x), \text{ where } x, y \in M_{n+1}$$

is an equivalence relation on M_{n+1} . Let $[x]_{R_{n+1}}$ denote the equivalence class of $x \in M_{n+1}$, obviously $\text{card } [x]_{R_{n+1}} \leq \text{card } Bij(M)$. Using the inequality $\text{card } M_{n+1} > \text{card } M_n \cdot \text{card } Bij(M)$ (valid both for finite and infinite M) we obtain that $\text{card } (M_{n+1}/R_{n+1}) > \text{card } M_n$. If $w \in M_{n+1}$ then $\forall g \in G \ g[w]_{R_{n+1}} = [w]_{R_{n+1}}$, hence the following G -set:

$$X_w := \left\{ \left\langle \underbrace{\{\dots\{x\}\dots\}}_{n-1 \text{ times}\{ \}}, [w]_{R_{n+1}} \right\rangle : x \in M_3 \right\}$$

is a subobject of M_{n+4} which is isomorphic to M_3 . Moreover, for every $s, t \in M_{n+1}$ if $[s]_{R_{n+1}} \neq [t]_{R_{n+1}}$ then $X_s \cap X_t = \emptyset$. This observation completes the proof.

References

- [1] N. Bourbaki, *Éléments de Mathématique, Livre II, Algèbre, Chapitre I, Structures algébriques*, Hermann, Paris 1958.
- [2] R. Goldblatt, *Topoi, the Categorical Analysis of Logic*, North-Holland, Amsterdam 1979.
- [3] P. T. Johnstone, *Topos Theory*, Academic Press, New York 1977.
- [4] B. Szociński, *Basic concepts of Klein geometries*, Zesz. Nauk. Politech. Śl. 1055, Mat.-Fiz. 62, 1990.
- [5] A. Tyszką, *On one combinatorial lemma and its geometric consequences*, Demonstr. Math. 25, 1992, pp. 579–582.
- [6] A. Tyszką, *On the notion a geometric object in a Klein space*, Rocznik Naukowo-Dydaktyczny Wyższej Szkoły Pedagogicznej w Krakowie 159, Prace Matematyczne XIII (Annales de l'Ecole Normale Supérieure à Cracovie 159, Travaux Mathématiques XIII), Kraków 1993, pp. 287–299.

Apoloniusz Tyszką
 Technical Faculty
 Hugo Kołłątaj University
 Balicka 116B, 30-149 Kraków, Poland
 E-mail address: rttyszka@cyf-kr.edu.pl